

## **Trade Name**                      **Forage, Feed, and Grazing Restrictions**

---

<b>2,4-D, others</b>	Do not forage or feed corn or sorghum fodder or graze animals on treated pasture areas for 7 days after application. Do not cut forage for hay within 30 days of application. Do not permit dairy animals or meat animals being finished for slaughter to forage treated forage grass fields within 3 days of slaughter or treated grain fields within 2 weeks of slaughter. Do not feed treated straw to livestock if a preharvest treatment or emergency treatment is applied. Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock. Refer to the label for other information.
<b>Aatrex</b>	Do not graze or feed forage from treated corn for 60 days after application, or illegal residues may result. Do not graze or feed forage from treated sweet corn for 45 days after application, or illegal residues may result. Do not graze or feed forage from treated sorghum for 60 days after a preemergence application or 45 days after a postemergence application, or illegal residues may result.
<b>Accent</b>	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of Accent application.
<b>Accent Gold WDG</b>	Do not harvest corn or graze or feed forage, hay or straw from treated areas to livestock within 85 days of Accent Gold application.
<b>Achieve</b>	Immature crops (forage) may be grazed or cut for hay 30 days after treatment. Mature straw and grain may be fed to livestock 45 days after treatment.
<b>Affinity BroadSpec</b>	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
<b>Affinity TankMix</b>	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
<b>Aim</b>	Do not feed treated soybean forage or soybean hay to livestock. Do not harvest small grains for forage within 7 days of application.
<b>alachlor, others</b>	Do not graze or harvest grain sorghum forage for 70 days following application. Do not feed forage, feed, or straw from treated soybeans. Do not ensile treated soybeans. Do not feed forage or hay from treated drybeans.
<b>Alphanex</b>	No restrictions on the label.
<b>Arrow</b>	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
<b>Assert</b>	Do not graze treated fields or cut treated forage for silage or hay. Wheat and barley straw may be fed or used for bedding.
<b>Assure II</b>	Do not graze treated fields or harvest for forage or hay.
<b>atrazine, others</b>	Do not graze or feed forage from treated corn for 60 days after application, or illegal residues may result. Do not graze or feed forage from treated sweet corn for 45 days after application, or illegal residues may result. Do not graze or feed forage from treated sorghum for 60 days after a preemergence application or 45 days after a postemergence application, or illegal residues may result.
<b>Avalanche</b>	No restrictions on the label.
<b>Axial</b>	Do not graze livestock or harvest forage for hay from treated areas for a minimum of 50 days following application.
<b>Balan</b>	None listed on the label.
<b>Banvel</b>	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest.
<b>Basagran</b>	Do not graze treated corn and sorghum fields for at least 12 days after treatment. Do not graze or cut treated soybean fields for forage or hay for at least 30 days after treatment.
<b>Basis</b>	Do not graze or feed forage, grain, or stover from treated areas to livestock within 30 days of application.
<b>Basis Gold</b>	Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of application.
<b>Beacon</b>	Do not graze or feed forage from Beacon-treated corn to livestock within 30 days after application. Do not harvest silage within 45 days after application.
<b>Betamix</b>	No restrictions on the label.

## **Trade Name**                      **Forage, Feed, and Grazing Restrictions**

---

<b>Betanex</b>	No restrictions on the label.
<b>Beyond</b>	Do not feed or graze wheat forage or hay until 30 days after application. Do not feed or graze treated canola.
<b>Bicep Lite II Magnum</b>	To avoid possible illegal residues, 1) do not graze treated or feed field corn forage from treated areas for 60 days following application.
<b>Bison</b>	Do not graze treated small grain fields within 45 days after application.
<b>Bison Advanced</b>	Do not graze treated small grain fields within 45 days after application.
<b>BnB Plus</b>	No restrictions on the label.
<b>Boundary 6.5</b>	Treated soybean plants may be grazed or fed to livestock 40 days after the last application.
<b>Brash</b>	For pasture areas, remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment. Do not graze or feed treated wheat prior to maturity.
<b>Bronate Advanced</b>	Do not graze treated small grain fields within 45 days after application.
<b>Buctril</b>	Do not cut corn for feed, fodder or graze within 45 days of application. Do not graze treated small grain fields for 45 days following treatment. Do not cut for feed or graze spring treated alfalfa (including underseeded with small grains) within 30 days following treatment. Do not cut for feed or graze fall treated alfalfa until spring.
<b>Buctril + atrazine</b>	Do not cut crop for feed or graze within 45 days after application.
<b>Bullet</b>	Do not graze treated area or feed treated forage to livestock for 21 days following application.
<b>Butoxone</b>	Do not graze or feed soybean hay within 60 days after application. Do not graze established alfalfa or feed straw or hay from treated established alfalfa to livestock within 30 days after application. Do not graze or feed seedling alfalfa, seedling clover, or seedling birdsfoot trefoil within 60 days after application.
<b>Butyrac</b>	Do not graze or feed soybean hay within 60 days after application. Do not graze established alfalfa or feed straw or hay from treated established alfalfa to livestock within 30 days after application. Do not graze or feed seedling alfalfa, seedling clover, or seedling birdsfoot trefoil within 60 days after application.
<b>Callisto</b>	Do not graze or feed treated plants for 45 days after treatment.
<b>Camix</b>	Do not graze or feed forage from treated areas for 45 days following application. Do not harvest forage or stover for 45 days following application.
<b>Celebrity Plus</b>	Do not apply within 32 days of forage harvest or 72 days of stover harvest.
<b>Charger MAX</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
<b>Charger MAX ATZ</b>	Do not graze treated or feed forage from treated areas for 30 days following application.
<b>Charger MAX ATZ Lite</b>	Do not graze treated or feed forage from treated areas for 30 days following application.
<b>Cimarron</b>	No grazing or haying restrictions.
<b>Cimarron Max</b>	Non-lactating animals: Remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Lactating animals: Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment.
<b>Cinch</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
<b>Cinch ATZ</b>	Do not graze treated or feed forage from treated areas for 30 days following application.
<b>Cinch ATZ Lite</b>	Do not graze treated or feed forage from treated areas for 30 days following application.

## **Trade Name**                      **Forage, Feed, and Grazing Restrictions**

---

<b>Clarity</b>	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest. SOYBEANS: Do not feed soybean fodder or hay following a preharvest application.
<b>Classic</b>	Do not graze treated fields or harvest for forage or hay.
<b>Clopyr AG</b>	Do not permit lactating dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields. Do not allow livestock to graze treated corn or harvest treated corn silage as feed within 40 days after last treatment. No restrictions on grazing or hay harvesting in rangeland or grass pasture situations. Do not feed or graze canola fields. Do not cut canola for forage. Do not graze sweet corn or popcorn fields for forage within 30 days of last treatment nor harvest for stover within 60 days of last treatment.
<b>Cobra</b>	Do not graze animals on green forage or stubble. Do not feed treated soybean silage to cattle. Do not utilize hay or straw for animal feed or bedding.
<b>Command</b>	Do not graze or harvest for food or feed cover crops planted less than 9 months after treatment.
<b>Confidence</b>	No restrictions.
<b>Confidence Xtra</b>	No restrictions on the label.
<b>Confidence Xtra 5.6L</b>	No restrictions on the label.
<b>Cornerstone</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>Crossbow</b>	Except for lactating dairy animals, there are no grazing restrictions following application. Lactating dairy animals: Do not allow lactating dairy animals to graze treated areas until the next growing season following application of this product. Do not harvest hay for 14 days after application. Grazed areas of non-crop land and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area. Slaughter restrictions: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.
<b>Curtail</b>	Do not graze lactating dairy cattle in treated pasture areas for 14 days after application. Remove meat animals from freshly treated areas 7 days before slaughter. Withdrawal is not needed if 2 weeks or more have elapsed since application. Do not cut treated grass for hay within 30 days after application. Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields.
<b>Curtail M</b>	Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment. Do not harvest hay from treated grain fields.
<b>Define</b>	Do not harvest corn forage (silage) within 75 days after a postemergence application. Do not graze or feed soybean forage, hay or straw to livestock.
<b>Degree</b>	No restrictions on the label.
<b>Degree Xtra</b>	No restrictions on the label.
<b>Des</b>	No restrictions on the label.
<b>Des-Phen-Etho</b>	No restrictions on the label.
<b>Discover</b>	Do not graze livestock or feed forage from treated areas for a minimum of 30 days following application. Do not feed hay for 30 days following application
<b>Discover NG</b>	Do not graze livestock or feed forage from treated areas for a minimum of 30 days following application. Do not feed hay for 30 days following application
<b>Distinct</b>	Do not graze or harvest hay from treated area.
<b>DP-Mix</b>	No restrictions on the label.
<b>Dual II Magnum</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.

## **Trade Name**                      **Forage, Feed, and Grazing Restrictions**

---

<b>Dual Magnum</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
<b>Durango</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not harvest alfalfa and feed livestock until a minimum of 36 hours has passed. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>Eptam</b>	Do not apply within 14 days of harvesting or grazing alfalfa. Do not apply within 45 days of harvesting or grazing ladino clover.
<b>Equip</b>	Do not graze or harvest corn forage within 45 days of application.
<b>Etho SC</b>	Do not graze livestock on treated crops.
<b>Ethotron SC</b>	Do not graze livestock on treated crops.
<b>Everest</b>	Treated wheat fields may be grazed at any time.
<b>Express XP</b>	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding or feed).
<b>Extreme</b>	Do not graze or feed treated soybean forage, hay or straw to livestock.
<b>Fargo</b>	Do not graze livestock on treated fields.
<b>Field Master</b>	Do not feed forage from corn or graze treated areas within 8 weeks after application.
<b>FirstRate</b>	Do not harvest soybeans for forage or hay for 14 days after application.
<b>Flexstar</b>	Do not graze treated areas or harvest for forage or hay.
<b>Fultime</b>	Do not apply this product within 60 days of harvest for forage use.
<b>Fusilade DX</b>	Do not graze or harvest for forage or hay.
<b>Fusion</b>	Do not graze or harvest for forage or hay.
<b>Garlon</b>	<p>Grazing or harvesting green forage: Following application of two quarts per acre or less: there are no restrictions for general livestock but do not graze or harvest green forage from treated area for 14 days after treatment for lactating dairy animals. Following applications of greater than two to six quarts per acre: do not graze or harvest green forage from treated area for 14 days after treatment for general livestock (unless total treated area is less than 25% of grazed area - in which case there is no grazing restriction) but do not graze or harvest green forage until the next season for lactating dairy animals.</p> <p>Haying Restrictions: Do not harvest hay to feed lactating dairy animals until the next growing season. For other livestock: Do not harvest hay for 7 days following application of two quarts per acre or less. Do not harvest hay for 14 days following application of greater than two up to 4 quarts per acre. Do not harvest for hay until next growing season for applications greater than 4 quarts per acre.</p> <p>Slaughter Restrictions: During the year following application, withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter.</p>
<b>Glyphomax XRT</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not harvest alfalfa and feed livestock until a minimum of 36 hours has passed. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>glyphosate, others</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>G-Max Lite</b>	Corn may be grazed or fed to livestock 40 or more days after application. Sorghum forage may be grazed or fed to livestock 60 or more days after application. Sorghum grain and fodder may be harvested and fed 80 or more days after application.
<b>Gramoxone</b>	Do not cut or harvest alfalfa within 42 days of application. Do not harvest treated sorghum for forage for 20 days following application. Do not graze or harvest soybeans for forage or hay before the R3 stage following preplant or preemergence applications. Do not graze or harvest soybean forage or hay following harvest aid applications. Do not graze treated sunflowers or feed treated sunflower forage to livestock following preharvest desiccation applications.

<b>Trade Name</b>	<b>Forage, Feed, and Grazing Restrictions</b>
-------------------	---

<b>Grazon P+D</b>	Do not allow lactating dairy animals to graze treated areas within 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days after application. Meat animals must be withdrawn from treated forage at least 3 days before slaughter. There are no other grazing restrictions for non-lactating dairy animals or other livestock.
<b>Harmony Extra XP</b>	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
<b>Harmony GT XP</b>	Do not graze or feed forage or hay from treated areas to livestock. (harvested straw may be used for bedding and/or feed).
<b>Harness</b>	No restrictions.
<b>Harness Xtra</b>	No restrictions on the label.
<b>Harness Xtra 5.6L</b>	No restrictions on the label.
<b>Hornet</b>	If field corn is grown for forage or ensiled, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.
<b>Intrro</b>	Do not feed forage or hay from treated drybeans. Do not feed treated soybean forage, feed or straw or ensile treated soybeans.
<b>Kerb</b>	Do not graze or harvest alfalfa for forage or dehydration west of the Mississippi River within 25 days (3.0 lbs/A Kerb 50W applied) or 45 days (3.0 to 4.0 lbs/A Kerb 50W applied). Do not graze or harvest clover, birdsfoot trefoil, crown vetch, sainfoin, or alfalfa for forage or dehydration east of the Mississippi River within 120 days.
<b>Keystone</b>	Do not apply within 60 days of harvest of field corn for forage uses.
<b>Keystone LA</b>	Do not apply within 60 days of harvest of field corn for forage uses.
<b>Laddock S-12</b>	Do not graze treated area or feed treated forage to livestock for 21 days after application.
<b>Lariat</b>	Do not graze treated area or feed treated forage to livestock for 21 days following application. Do not graze or harvest treated grain sorghum for 70 days following application.
<b>Lasso</b>	Do not graze or harvest forage of treated grain sorghum for 70 days following application. Do not feed treated soybean forage, feed or straw or ensile treated soybeans.
<b>Liberty</b>	Do not apply within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder. Do not graze treated canola crop or cut for hay.
<b>Lightning</b>	There should be an interval of at least 45 days between application and corn harvest (silage, fodder, or grain). Do not graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after application.
<b>Lorox</b>	Do not graze or feed sorghum forage or silage from treated fields to dairy animals. Do not graze or feed soybean forage from treated areas to livestock.
<b>Lumax</b>	Do not graze or feed forage from treated areas for 45 days following application. Do not harvest forage or stover for 45 days following application.
<b>Marksman</b>	Corn can be harvested or grazed for feed once the crop has reached the ensilage (milk stage) or later in maturity. Sorghum may be harvested or grazed for feed once the crop has reached the mature grain stage.
<b>Matrix</b>	No restrictions on the label.
<b>MCPA, others</b>	Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.
<b>Micro Tech</b>	Do not graze or harvest forage of treated grain sorghum for 70 days following application. Do not feed treated soybean forage, feed or straw or ensile treated soybeans.
<b>Milestone</b>	There are no restrictions on grazing or hay harvest following application of Milestone at labeled rates. Do not transfer grazing animals from areas treated with Milestone to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
<b>Moxy</b>	Do not cut corn for feed, fodder or graze within 45 days of application. Do not graze treated small grain fields for 45 days following treatment. Do not cut for feed or graze spring treated alfalfa (including underseeded with small grains) within 30 days following treatment. Do not cut for feed or graze fall treated alfalfa until spring at least 60 days after treatment.

<b>Trade Name</b>	<b>Forage, Feed, and Grazing Restrictions</b>
-------------------	---

<b>Northstar</b>	Do not graze or feed forage from treated corn to livestock within 30 days after application. Do not harvest silage within 45 days after application.
<b>Nortron SC</b>	Do not graze livestock on treated crops.
<b>Option</b>	Do not graze or harvest corn forage within 45 days after Option treatment.
<b>Outlook</b>	Corn may be grazed or fed to livestock 40 days after application. Do not graze or feed soybean forage, hay, or straw to livestock. Sorghum forage may be grazed or fed to livestock 60 days after application. Sorghum grain or fodder may be harvested and fed 80 days after application.
<b>Overdrive</b>	Pasture and rangeland grass can be grazed or harvested for livestock feed immediately after application.
<b>Parallel</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application. Do not graze or feed treated soybean forage, hay, or straw to livestock.
<b>Parallel PCS</b>	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application. Do not graze or feed treated soybean forage, hay, or straw to livestock.
<b>Pendant</b>	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
<b>Pendimax</b>	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
<b>Permit</b>	Allow 30 days from application before grazing domestic livestock, harvesting forage, or harvesting silage.
<b>Phen-Des 8+8</b>	No restrictions on the label.
<b>Phoenix</b>	Do not graze animals on green forage or stubble. Do not feed treated soybean silage to cattle. Do not utilize hay or straw for animal feed or bedding.
<b>Plateau</b>	Do not harvest treated areas for hay within 7 days of treatment.
<b>Poast</b>	For alfalfa, birdsfoot trefoil, and sainfoin do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 14 days of cutting for dry hay. For clover do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 20 days of cutting for dry hay. Do not harvest or graze other cover crops. For Poast Protected field corn, do not apply within 60 days of harvesting for grain or fodder or within 45 days of harvesting for forage or silage. Only processed soybean meal from seed or soybean hay may be fed to animals.
<b>Poast Plus</b>	For alfalfa, birdsfoot trefoil, and sainfoin do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 14 days of cutting for dry hay. For clover do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 20 days of cutting for dry hay. Do not harvest or graze other cover crops. For Poast Protected field corn, do not apply within 60 days of harvesting for grain or fodder or within 45 days of harvesting for forage or silage. Only processed soybean meal from seed or soybean hay may be fed to animals.
<b>Princep</b>	Do not graze treated areas.
<b>Prism</b>	Do not graze treated soybeans or feed treated forage or hay to livestock.
<b>Progress</b>	No restrictions on the label.
<b>Propel</b>	Corn may be grazed or fed to livestock 40 days after application. Do not graze or feed soybean forage, hay, or straw to livestock. Sorghum forage may be grazed or fed to livestock 60 days after application. Sorghum grain or fodder may be harvested and fed 80 days after application.
<b>Prowl</b>	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
<b>Prowl H20</b>	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.

<b>Trade Name</b>	<b>Forage, Feed, and Grazing Restrictions</b>
-------------------	---

<b>Puma</b>	No restrictions on the label.
<b>Pursuit</b>	Do not graze or feed treated soybean forage, hay or straw to livestock. Do not graze or feed Clearfield corn forage, silage, fodder, or grain for at least 45 days after application. Do not feed or graze alfalfa for 30 days following application.
<b>Pursuit Plus</b>	Do not graze or feed Clearfield corn forage, silage, fodder, or grain for at least 45 days after application. Do not graze or feed treated soybean forage, hay or straw to livestock.
<b>Python</b>	Do not graze or feed soybean forage, hay or straw to livestock.
<b>Raptor</b>	Do not graze or feed treated soybean forage, hay or straw to livestock. No waiting period between application of Raptor and cutting or feeding alfalfa forage or hay.
<b>Rave</b>	Except for lactating dairy animals, there are no grazing restrictions following an application of Rave. Treated areas should not be grazed by lactating dairy animals before 7 days after treatment. Animals cannot be removed from treated areas for slaughter less than 30 days after application. Do not harvest hay from treated fields for 37 days after treatment.
<b>Redeem</b>	GRAZING or HARVESTING GREEN FORAGE: For lactating dairy animals, do not graze or harvest green forage from treated area for 14 days after treatment. For other livestock, there are no grazing restrictions. HARVESTING DRY FORAGE: For lactating dairy animals, do not harvest hay until the next growing season. For other livestock, do not harvest hay for 7 days after treatment. SLAUGHTER RESTRICTIONS: Withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter. This restriction applies to grazing during the season of treatment or hay harvested during the season of treatment.
<b>Reflex</b>	Do not graze treated areas or harvest for forage or hay. Do not graze rotated small grain crops or harvest forage or straw for livestock.
<b>Remedy</b>	GRAZING of HARVESTING GREEN FORAGE: For lactating dairy animals, do not graze or harvest green forage for 14 days (if 2 quarts/acre or less is applied) or until the next season (greater than 2 up to 6 quarts/acre product applied). For other livestock, no grazing restrictions (if 2 quarts/acre or less is applied or less than 25% of the grazing area is treated at a rate greater than 2 quarts/acre), otherwise do not graze or harvest green forage for 14 days after treatment (greater than 2 up to 6 quarts/acre product applied). HARVESTING DRY FORAGE: Do not harvest hay for lactating dairy animals until the next year. For other animals, do not harvest for hay until 7 days after treatment (2 quarts/acre or less is applied) or 14 days after treatment (greater than 2 up to 4 quarts/acre product applied). If greater than 4 quarts/acre is applied, do not harvest for hay until the next growing season. SLAUGHTER RESTRICTIONS: Withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter. This restriction applies to grazing during the season following treatment or hay harvested during the season following treatment.
<b>Resolve</b>	Do not graze, feed forage, grain or fodder (stover) from treated areas to livestock within 30 days of application.
<b>Resource</b>	Do not graze animals on green corn forage or use for seed less than 28 days after application. Do not graze treated soybean fields or harvest for forage or hay.
<b>Rezult</b>	Do not graze treated soybean fields and do not feed treated soybean forage (green succulent) or ensilage to livestock. Do not cut treated soybean fields for forage or hay for at least 30 days after treatment. Do not graze treated Poast Protected corn for at least 12 days after treatment.
<b>Rimfire</b>	Forage = 30 days PHI, Grain and Straw = 71 days PHI.
<b>Ro-Neet</b>	No restrictions on the label.
<b>Roundup</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>Roundup Original MAX</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>Roundup UltraMAX II</b>	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.

## **Trade Name**                      **Forage, Feed, and Grazing Restrictions**

---

<b>Roundup WeatherMAX</b>	For preharvest applications in corn, allow at least 7 days between application and feeding of stover or grain. Do not feed or graze treated grain sorghum fodder or ensile treated grain sorghum. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. If the preharvest application rate in soybeans is less than 22oz/A the grazing restriction is reduced to 14 days. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
<b>Saber</b>	Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied.
<b>Salvo</b>	Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied. Do not forage or feed corn fodder for 7 days following application.
<b>Section</b>	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
<b>Select</b>	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
<b>Select Max</b>	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
<b>Sencor</b>	Treated soybean vines may be fed to livestock 40 days after application. Field corn treated with Sencor may be grazed or harvested for silage or grain 60 days after treatment. Do not graze or harvest alfalfa within 28 days after application.
<b>Shotgun</b>	Do not graze or feed forage from treated areas for 21 days following application.
<b>Silverado</b>	Do not apply Silverado within 30 days of harvesting wheat forage, 50 days for hay, and 55 days for grain or straw.
<b>Sinbar</b>	No restrictions on the label.
<b>sodium chlorate, others</b>	
<b>Sonalan</b>	Do not graze or forage crop grown in treated soil or cut for hay or silage.
<b>Spartan</b>	Do not feed treated soybean forage or soybean hay to livestock.
<b>Spike</b>	No restrictions on the label.
<b>Stampede</b>	Do not graze treated crop or cut for green chop feed.
<b>Starane</b>	Do not allow livestock to graze treated areas or harvest treated forage within 47 days of application in corn, 40 days in grain sorghum, or 7 days in oats, barley or wheat. Do not apply less than 90 days before harvest of grain or stover in corn, 70 days in grain sorghum, or 40 days in oats, barley or wheat.
<b>Starane + Saber</b>	Do not allow livestock to graze treated areas or harvest treated forage within 14 days in oats, barley or wheat. Do not apply closer than 14 days before cutting of hay or harvesting of straw.
<b>Starane + Salvo</b>	Do not allow livestock to graze treated areas or harvest treated forage within 14 days in oats, barley or wheat. Do not apply closer than 14 days before cutting of hay or harvesting of straw.
<b>Starane + Sword</b>	Do not allow livestock to graze treated areas or harvest treated forage within 7 days in oats, barley or wheat. Do not apply closer than 40 days before cutting of hay or harvesting of straw.
<b>Steadfast</b>	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of application.
<b>Steadfast ATZ</b>	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of application.
<b>Sterling</b>	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest.
<b>Sterling Plus</b>	Corn can be harvested or grazed for feed once the crop has reached the ensilage (milk stage) or later in maturity. Sorghum may be harvested or grazed for feed once the crop has reached the mature grain stage.

<b>Trade Name</b>	<b>Forage, Feed, and Grazing Restrictions</b>
-------------------	---

<b>Stinger</b>	Do not permit lactating dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields. Do not allow livestock to graze treated corn or harvest treated corn silage as feed within 40 days after last treatment. No restrictions on grazing or hay harvesting in rangeland or grass pasture situations. Do not feed or graze canola fields. Do not cut canola for forage. Do not graze sweet corn or popcorn fields for forage within 30 days of last treatment nor harvest for stover within 60 days of last treatment.
<b>Storm</b>	Do not use treated plants for feed or forage.
<b>Surpass</b>	No restrictions on the label.
<b>Sword</b>	Do not forage or graze dairy and meat animals on treated areas within seven days of slaughter.
<b>Synchrony XP</b>	Do not graze treated fields or harvest for forage or hay.
<b>Targa</b>	Do not graze livestock in treated areas. In addition, do not feed forage, hay, or straw from treated areas to livestock. Do not feed sugarbeet tops within 60 days of last application.
<b>Topnotch</b>	No restrictions on the label.
<b>Tordon 22K</b>	When applying more than 0.5 lb ai/A to grass pastures, do not cut grass for feed within two weeks after treatment. Meat animals grazing for up to two weeks after treatment should be removed from treated areas three days prior to slaughter. Do not graze lactating dairy animals on treated areas within two weeks after treatment. Do not graze or feed small grain forage from treated areas for 2 weeks after treatment. Do not harvest hay from treated grain fields.
<b>Touchdown</b>	Do not graze or feed corn forage or fodder following hooded sprayer applications. Allow a minimum of 50 days between postemergence application and harvest of forage of glyphosate-tolerant corn. Allow at least 25 days before grazing or harvesting for livestock feed following harvest aid applications in conventional soybeans. Do not graze or harvest treated glyphosate-tolerant soybeans for forage or hay. Do not graze or feed sunflower forage.
<b>Treflan</b>	Do not cut or graze alfalfa within 21 days after application.
<b>trifluralin, others</b>	Do not cut or graze alfalfa within 21 days after application.
<b>Trust</b>	Do not cut or graze alfalfa within 21 days after application.
<b>Ultra Blazer</b>	Do not use treated plants for feed or forage.
<b>UpBeet</b>	No restrictions on grazing or feeding of crop residue to livestock.
<b>Valor</b>	Do not graze treated fields or feed treated forage or hay to livestock.
<b>Velpar</b>	Do not graze or feed forage or hay to livestock within 30 days following application.
<b>Velpar AlfaMax MP</b>	Do not graze or feed forage to livestock within 30 days following application.
<b>Weedmaster</b>	Remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment.
<b>WideMatch</b>	Do not allow livestock to graze treated areas or harvest treated small grain forage within 7 days of application. Do not apply closer than 14 days before cutting small grains for hay or 40 days before harvesting of straw. Do not allow livestock to graze treated areas or harvest treated corn forage within 47 days of application. Do not apply less than 90 days before harvest of corn stover. Do not allow livestock to graze treated areas or harvest treated sweet corn forage within 31 days of application. Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.
<b>Yukon</b>	Corn can be grazed or harvested for feed after the crop reaches the ensilage (milk) stage at least 30 days after application.