

Trade Name **Forage, Feed, and Grazing Restrictions**

2,4-D, generic	Do not forage or feed corn or sorghum fodder or graze animals on treated pasture areas for 7 days after application. Do not cut forage for hay within 30 days of application. Do not permit dairy animals or meat animals being finished for slaughter to forage treated forage grass fields within 3 days of slaughter or treated grain fields within 2 weeks of slaughter. Do not feed treated straw to livestock if a preharvest treatment or emergency treatment is applied. Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock. Refer to the label for other information.
Aatrex	Do not graze or feed forage from treated corn for 60 days after application, or illegal residues may result. Do not graze or feed forage from treated sweet corn for 45 days after application, or illegal residues may result. Do not graze or feed forage from treated sorghum for 60 days after a preemergence application or 45 days after a postemergence application, or illegal residues may result.
Accent	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of Accent application.
Achieve	Immature crops (forage) may be grazed or cut for hay 30 days after treatment. Mature straw and grain may be fed to livestock 45 days after treatment.
Affinity BroadSpec	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
Affinity TankMix	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
Aim	Do not feed treated soybean forage or soybean hay to livestock. Do not harvest small grains for forage within 7 days of application.
alachlor, generic	Do not graze or harvest grain sorghum forage for 70 days following application. Do not feed forage, feed, or straw from treated soybeans. Do not ensile treated soybeans. Do not feed forage or hay from treated drybeans.
Alphanex	No restrictions on the label.
Arrow	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
Assert	Do not graze treated fields or cut treated forage for silage or hay. Wheat and barley straw may be fed or used for bedding.
Assure II	Do not graze treated fields or harvest for forage or hay.
atrazine, generic	Do not graze or feed forage from treated corn for 60 days after application, or illegal residues may result. Do not graze or feed forage from treated sweet corn for 45 days after application, or illegal residues may result. Do not graze or feed forage from treated sorghum for 60 days after a preemergence application or 45 days after a postemergence application, or illegal residues may result.
Authority First	Do not feed treated soybean forage or soybean hay to livestock.
Authority MTZ	Do not graze treated fields or harvest for forage or hay.
Axial	Do not graze livestock or harvest forage for hay from treated areas for a minimum of 50 days following application.
Balan	None listed on the label.
Banvel	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest.
Basagran	Do not graze treated corn and sorghum fields for at least 12 days after treatment. Do not graze or cut treated soybean fields for forage or hay for at least 30 days after treatment.
Basis	Do not graze or feed forage, grain, or stover from treated areas to livestock within 30 days of application.
Beacon	Do not graze or feed forage from Beacon-treated corn to livestock within 30 days after application. Do not harvest silage within 45 days after application.
Betamix	No restrictions on the label.
Betanex	No restrictions on the label.
Beyond	Do not feed or graze wheat forage or hay until 30 days after application. Do not feed or graze treated canola.

Trade Name **Forage, Feed, and Grazing Restrictions**

Bicep Lite II Magnum	To avoid possible illegal residues, 1) do not graze treated or feed field corn forage from treated areas for 60 days following application.
Bison	Do not graze treated small grain fields within 45 days after application.
Bison Advanced	Do not graze treated small grain fields within 45 days after application.
BnB Plus	No restrictions on the label.
Boundary 6.5	Treated soybean plants may be grazed or fed to livestock 40 days after the last application.
Brash	For pasture areas, remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment. Do not graze or feed treated wheat prior to maturity.
Breakfree	No restrictions on the label.
Breakfree ATZ	Do not harvest field corn for forage uses within 60 days of application.
Breakfree ATZ Lite	Do not harvest field corn for forage uses within 60 days of application.
Broclean	Do not cut corn for feed, fodder or graze within 45 days of application. Do not graze treated small grain fields for 45 days following treatment. Do not cut for feed or graze spring treated alfalfa (including underseeded with small grains) within 30 days following treatment. Do not cut for feed or graze fall treated alfalfa until spring.
Bromac	Do not graze treated small grain fields within 45 days after application.
Bromac Advanced	Do not graze treated small grain fields within 45 days after application.
Bronate Advanced	Do not graze treated small grain fields within 45 days after application.
Brozine	Do not cut crop for feed or graze within 45 days after application.
Buctril	Do not cut corn for feed, fodder or graze within 45 days of application. Do not graze treated small grain fields for 45 days following treatment. Do not cut for feed or graze spring treated alfalfa (including underseeded with small grains) within 30 days following treatment. Do not cut for feed or graze fall treated alfalfa until spring.
Buctril + atrazine	Do not cut crop for feed or graze within 45 days after application.
Bullet	Do not graze treated area or feed treated forage to livestock for 21 days following application.
Butoxone	Do not graze or feed soybean hay within 60 days after application. Do not graze established alfalfa or feed straw or hay from treated established alfalfa to livestock within 30 days after application. Do not graze or feed seedling alfalfa, seedling clover, or seedling birdsfoot trefoil within 60 days after application.
Butyrac	Do not graze or feed soybean hay within 60 days after application. Do not graze established alfalfa or feed straw or hay from treated established alfalfa to livestock within 30 days after application. Do not graze or feed seedling alfalfa, seedling clover, or seedling birdsfoot trefoil within 60 days after application.
Cadence	No restrictions on the label.
Cadence ATZ	Do not harvest field corn for forage uses within 60 days of application.
Cadence ATZ Lite	Do not harvest field corn for forage uses within 60 days of application.
Callisto	Do not graze or feed treated plants for 45 days after treatment.
Camix	Do not graze or feed forage from treated areas for 45 days following application. Do not harvest forage or stover for 45 days following application.
Charger Basic	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
Charger MAX	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
Charger MAX ATZ	Do not graze treated or feed forage from treated areas for 30 days following application.
Charger MAX ATZ Lite	Do not graze treated or feed forage from treated areas for 30 days following application.
Cimarron	No grazing or haying restrictions.

Trade Name **Forage, Feed, and Grazing Restrictions**

Cimarron Max	Non-lactating animals: Remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Lactating animals: Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment.
Cinch	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
Cinch ATZ	Do not graze treated or feed forage from treated areas for 30 days following application.
Cinch ATZ Lite	Do not graze treated or feed forage from treated areas for 30 days following application.
Clarity	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest. SOYBEANS: Do not feed soybean fodder or hay following a preharvest application.
Classic	Do not graze treated fields or harvest for forage or hay.
Clearmax	Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.
Clopyr AG	Do not permit lactating dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields. Do not allow livestock to graze treated corn or harvest treated corn silage as feed within 40 days after last treatment. No restrictions on grazing or hay harvesting in rangeland or grass pasture situations. Do not feed or graze canola fields. Do not cut canola for forage. Do not graze sweet corn or popcorn fields for forage within 30 days of last treatment nor harvest for stover within 60 days of last treatment.
Cobra	Do not graze animals on green forage or stubble. Do not feed treated soybean silage to cattle. Do not utilize hay or straw for animal feed or bedding.
Colt AS	Do not allow livestock to graze treated areas or harvest treated small grain forage within 7 days of application. Do not apply closer than 14 days before cutting small grains for hay or 40 days before harvesting of straw. Do not allow livestock to graze treated areas or harvest treated corn forage within 47 days of application. Do not apply less than 90 days before harvest of corn stover. Do not allow livestock to graze treated areas or harvest treated sweet corn forage within 31 days of application. Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.
Command	Do not graze or harvest for food or feed cover crops planted less than 9 months after treatment.
Confidence	No restrictions on the label.
Confidence Xtra	No restrictions on the label.
Confidence Xtra 5.6L	No restrictions on the label.
Cornerstone	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
Cornerstone Plus	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
Crossbow	Except for lactating dairy animals, there are no grazing restrictions following application. Lactating dairy animals: Do not allow lactating dairy animals to graze treated areas until the next growing season following application of this product. Do not harvest hay for 14 days after application. Grazed areas of non-crop land and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area. Slaughter restrictions: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

Trade Name **Forage, Feed, and Grazing Restrictions**

Curtail	Do not graze lactating dairy cattle in treated pasture areas for 14 days after application. Remove meat animals from freshly treated areas 7 days before slaughter. Withdrawal is not needed if 2 weeks or more have elapsed since application. Do not cut treated grass for hay within 30 days after application. Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields.
Curtail M	Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment. Do not harvest hay from treated grain fields.
Define	Do not harvest corn forage (silage) within 75 days after a postemergence application. Do not graze or feed soybean forage, hay or straw to livestock.
Degree	No restrictions on the label.
Degree Xtra	No restrictions on the label.
Des	No restrictions on the label.
Des-Phen-Etho	No restrictions on the label.
Dimetric	Treated soybean vines may be fed to livestock 40 days after application. Field corn treated with Sencor may be grazed or harvested for silage or grain 60 days after treatment. Do not graze or harvest alfalfa within 28 days after application.
Discover	Do not graze livestock or feed forage from treated areas for a minimum of 30 days following application. Do not feed hay for 30 days following application
Discover NG	Do not graze livestock or feed forage from treated areas for a minimum of 30 days following application. Do not feed hay for 30 days following application
Distinct	Do not graze or harvest hay from treated area.
Double Up B+D	Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 45 days after treatment.
DP-Mix	No restrictions on the label.
Dual II Magnum	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
Dual Magnum	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application.
Duramax	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not harvest alfalfa and feed livestock until a minimum of 36 hours has passed. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
Durango DMA	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not harvest alfalfa and feed livestock until a minimum of 36 hours has passed. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
Enlite	Do not graze treated fields or harvest for forage or hay.
Eptam	Do not apply within 14 days of harvesting or grazing alfalfa. Do not apply within 45 days of harvesting or grazing ladino clover.
Equip	Do not graze or harvest corn forage within 45 days of application.
Eradicane	No restrictions on the label.
Etho SC	Do not graze livestock on treated crops.
Ethotron SC	Do not graze livestock on treated crops.
Everest	Treated wheat fields may be grazed at any time.
Express	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding or feed).
Extreme	Do not graze or feed treated soybean forage, hay or straw to livestock.

Trade Name	Forage, Feed, and Grazing Restrictions
-------------------	---

Fargo	Do not graze livestock on treated fields.
Field Master	Do not feed forage from corn or graze treated areas within 8 weeks after application.
FirstRate	Do not harvest soybeans for forage or hay for 14 days after application.
Flexstar	Do not graze treated areas or harvest for forage or hay.
ForeFront R&P	Do not harvest forage for hay within 7 days of application. Do not transfer grazing animals from areas treated with ForeFront R&P to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
Fultime	Do not apply this product within 60 days of harvest for forage use.
Fusilade DX	Do not graze or harvest for forage or hay.
Fusion	Do not graze or harvest for forage or hay.
Gangster	Do not graze treated fields or feed treated forage or hay to livestock.
Garlon	<p>Grazing or harvesting green forage: Following application of two quarts per acre or less: there are no restrictions for general livestock but do not graze or harvest green forage from treated area for 14 days after treatment for lactating dairy animals. Following applications of greater than two to six quarts per acre: do not graze or harvest green forage from treated area for 14 days after treatment for general livestock (unless total treated area is less than 25% of grazed area - in which case there is no grazing restriction) but do not graze or harvest green forage until the next season for lactating dairy animals.</p> <p>Haying Restrictions: Do not harvest hay to feed lactating dairy animals until the next growing season. For other livestock: Do not harvest hay for 7 days following application of two quarts per acre or less. Do not harvest hay for 14 days following application of greater than two up to 4 quarts per acre. Do not harvest for hay until next growing season for applications greater than 4 quarts per acre.</p> <p>Slaughter Restrictions: During the year following application, withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter.</p>
glyphosate, generic	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
G-Max Lite	Corn may be grazed or fed to livestock 40 or more days after application. Sorghum forage may be grazed or fed to livestock 60 or more days after application. Sorghum grain and fodder may be harvested and fed 80 or more days after application.
Gramoxone	Do not cut or harvest alfalfa within 42 days of application. Do not harvest treated sorghum for forage for 20 days following application. Do not graze or harvest soybeans for forage or hay before the R3 stage following preplant or preemergence applications. Do not graze or harvest soybean forage or hay following harvest aid applications. Do not graze treated sunflowers or feed treated sunflower forage to livestock following preharvest desiccation applications.
Grazon P+D	Do not allow lactating dairy animals to graze treated areas within 7 days after application. Do not harvest grass cut for hay from treated areas for 30 days after application. Meat animals must be withdrawn from treated forage at least 3 days before slaughter. There are no other grazing restrictions for non-lactating dairy animals or other livestock.
Guardsman Max	Field corn forage may be grazed or fed to livestock 60 days or more after application. Sweet corn forage may be grazed or fed to livestock 60 days or more after application. Sorghum forage may be grazed or fed to livestock 60 days or more after a preemergence application. Sorghum forage may be grazed or fed to livestock 45 days or more after a postemergence application. Sorghum grain or fodder may be harvested and fed 80 days or more after application.
Halex GT	Do not graze or feed forage from treated areas for 45 days following application.
Harmony	Do not graze or feed forage or hay from treated areas to livestock. (harvested straw may be used for bedding and/or feed).
Harmony Extra	Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (harvested straw may be used for bedding and/or feed).
Harness	No restrictions.
Harness Xtra	No restrictions on the label.

Trade Name **Forage, Feed, and Grazing Restrictions**

Harness Xtra 5.6L	No restrictions on the label.
Hornet WDG	If field corn is grown for forage or ensiled, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.
Huskie	Do not graze or harvest forage within 25 days, grain and straw within 60 days after application.
Impact	Do not graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after an application.
Imperium	No restrictions on the label.
Intensity	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
Intensity One	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
INTRRO	Do not feed forage or hay from treated drybeans. Do not feed treated soybean forage, feed or straw or ensile treated soybeans.
Kerb	Do not graze or harvest alfalfa for forage or dehydration west of the Mississippi River within 25 days (3.0 lbs/A Kerb 50W applied) or 45 days (3.0 to 4.0 lbs/A Kerb 50W applied). Do not graze or harvest clover, birdsfoot trefoil, crown vetch, sainfoin, or alfalfa for forage or dehydration east of the Mississippi River within 120 days.
Keystone	Do not apply within 60 days of harvest of field corn for forage uses.
Keystone LA	Do not apply within 60 days of harvest of field corn for forage uses.
Laddok S-12	Do not graze treated area or feed treated forage to livestock for 21 days after application.
Lariat	Do not graze treated area or feed treated forage to livestock for 21 days following application. Do not graze or harvest treated grain sorghum for 70 days following application.
Laudis	Do not graze livestock or harvest corn forage within 45 days of application.
Liberty	Do not apply within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder. Do not graze treated canola crop or cut for hay.
Lumax	Do not graze or feed forage from treated areas for 45 days following application. Do not harvest forage or stover for 45 days following application.
Makaze	Do not graze or feed corn forage or fodder following applications through hooded sprayers. Do not feed or graze treated milo fodder or ensile treated milo. Do not graze or harvest treated soybeans within 25 days of the last preharvest application. Do not feed or graze sunflower following application. For any other broadcast postemergence treatment, do not harvest or feed treated vegetation for 8 weeks following application.
Marksman	Corn can be harvested or grazed for feed once the crop has reached the ensilage (milk stage) or later in maturity. Sorghum may be harvested or grazed for feed once the crop has reached the mature grain stage.
MCPA, generic	Do not forage or graze meat animals on treated areas within 7 days of slaughter. Do not forage or graze dairy animals on treated areas within 7 days after treatment.
Micro Tech	Do not graze or harvest forage of treated grain sorghum for 70 days following application. Do not feed treated soybean forage, feed or straw or ensile treated soybeans.
Milestone	There are no restrictions on grazing or hay harvest following application of Milestone at labeled rates. Do not transfer grazing animals from areas treated with Milestone to areas where sensitive broadleaf crops occur without first allowing 3 days of grazing on an untreated pasture. Otherwise, urine and manure may contain enough aminopyralid to cause injury to sensitive broadleaf plants.
Moxy	Do not cut corn for feed, fodder or graze within 45 days of application. Do not graze treated small grain fields for 45 days following treatment. Do not cut for feed or graze spring treated alfalfa (including underseeded with small grains) within 30 days following treatment. Do not cut for feed or graze fall treated alfalfa until spring at least 60 days after treatment.
Northstar	Do not graze or feed forage from treated corn to livestock within 30 days after application. Do not harvest silage within 45 days after application.
Nortron SC	Do not graze livestock on treated crops.
Option	Do not graze or harvest corn forage within 45 days after Option treatment.
Orion	Livestock may be grazed on treated crops 7 days following application.

Trade Name **Forage, Feed, and Grazing Restrictions**

Outlook	Corn may be grazed or fed to livestock 40 days after application. Do not graze or feed soybean forage, hay, or straw to livestock. Sorghum forage may be grazed or fed to livestock 60 days after application. Sorghum grain or fodder may be harvested and fed 80 days after application.
Overdrive	Pasture and rangeland grass can be grazed or harvested for livestock feed immediately after application.
Parallel	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application. Do not graze or feed treated soybean forage, hay, or straw to livestock.
Parallel PCS	Do not graze or feed forage from treated corn for 30 days following application. Do not cut pod crops (drybeans, lupines, or peas) for hay within 120 days following application. Do not graze or feed treated soybean forage, hay, or straw to livestock.
Parallel Plus	Do not graze or feed forage from treated areas for 60 days following application for all types of corn except sweet corn. For sweet corn, do not graze or feed forage from treated areas for 45 days following application. Do not graze or feed forage from preemergent treated sorghum for 60 days. Do not graze or feed forage from post-emergent treated sorghum for 45 days.
Pendant	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
Pendimax	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
Permit	Allow 30 days from application before grazing domestic livestock, harvesting forage, or harvesting silage.
Phen-Des 8+8	No restrictions on the label.
Phoenix	Do not graze animals on green forage or stubble. Do not feed treated soybean silage to cattle. Do not utilize hay or straw for animal feed or bedding.
Plateau	Do not harvest treated areas for hay within 7 days of treatment.
Poast	For alfalfa, birdsfoot trefoil, and sainfoin do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 14 days of cutting for dry hay. For clover do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 20 days of cutting for dry hay. Do not harvest or graze other cover crops. For Poast Protected field corn, do not apply within 60 days of harvesting for grain or fodder or within 45 days of harvesting for forage or silage. Only processed soybean meal from seed or soybean hay may be fed to animals.
Poast Plus	For alfalfa, birdsfoot trefoil, and sainfoin do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 14 days of cutting for dry hay. For clover do not apply within 7 days of grazing, feeding, or cutting for undried forage or within 20 days of cutting for dry hay. Do not harvest or graze other cover crops. For Poast Protected field corn, do not apply within 60 days of harvesting for grain or fodder or within 45 days of harvesting for forage or silage. Only processed soybean meal from seed or soybean hay may be fed to animals.
Prefix	Do not graze treated areas or harvest for forage or hay. Do not graze rotated small grain crops or harvest forage or straw for livestock.
Princep	Do not graze treated areas.
Progress	No restrictions on the label.
Propel	Corn may be grazed or fed to livestock 40 days after application. Do not graze or feed soybean forage, hay, or straw to livestock. Sorghum forage may be grazed or fed to livestock 60 days after application. Sorghum grain or fodder may be harvested and fed 80 days after application.
Propel ATZ	Field corn may be grazed or fed to livestock 60 or more days after application. Sweet corn may be grazed or fed to livestock 45 or more days after application.
Propel ATZ Lite	Field corn may be grazed or fed to livestock 60 or more days after application. Sweet corn may be grazed or fed to livestock 45 or more days after application.
Prowl	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.

Trade Name**Forage, Feed, and Grazing Restrictions**

Prowl H20	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields. Do not apply within 60 days of harvesting wheat straw, within 28 days of harvesting wheat hay, or within 11 days of harvesting wheat forage. Do not apply within 50 days of harvesting alfalfa for forage or hay.
Puma	No restrictions on the label.
Pursuit	Do not graze or feed treated soybean forage, hay or straw to livestock. Do not graze or feed Clearfield corn forage, silage, fodder, or grain for at least 45 days after application. Do not feed or graze alfalfa for 30 days following application.
Pursuit Plus	Do not graze or feed Clearfield corn forage, silage, fodder, or grain for at least 45 days after application. Do not graze or feed treated soybean forage, hay or straw to livestock.
Python	Do not graze or feed soybean forage, hay or straw to livestock.
Rage D-Tech	Do not forage or feed corn fodder for 7 days following application. Do not graze dairy animals for 7 days following application on grass pastures or rangelands. Do not harvest grass for hay for 30 days following application. Remove meat animals from treated pastures or rangeland 3 days before slaughter. Do not feed sorghum fodder for 7 days following application. Do not graze meat animals on treated sorghum within 3 days of slaughter. Do not graze dairy animals on treated sorghum within 7 days after application. Do not graze dairy animals or meat animals being finished for slaughter for 14 days following application on treated small grains. Do not feed treated small grain straw to livestock.
Raptor	Do not graze or feed treated soybean forage, hay or straw to livestock. No waiting period between application of Raptor and cutting or feeding alfalfa forage or hay.
Reflex	Do not graze treated areas or harvest for forage or hay. Do not graze rotated small grain crops or harvest forage or straw for livestock.
Remedy Ultra	Do not allow lactating dairy animals to graze treated areas until the next growing season following application. Do not harvest hay for lactating dairy animals for 14 days after application. SLAUGHTER RESTRICTIONS: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.
Resolve	Do not graze, feed forage, grain or fodder (stover) from treated areas to livestock within 30 days of application.
Resource	Do not graze animals on green corn forage or use for seed less than 28 days after application. Do not graze treated soybean fields or harvest for forage or hay.
Rezult	Do not graze treated soybean fields and do not feed treated soybean forage (green succulent) or ensilage to livestock. Do not cut treated soybean fields for forage or hay for at least 30 days after treatment. Do not graze treated Poast Protected corn for at least 12 days after treatment.
Rifle	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest.
Rifle Plus	Corn can be harvested or grazed for feed once the crop has reached the ensilage (milk stage) or later in maturity. Sorghum may be harvested or grazed for feed once the crop has reached the mature grain stage.
Rifle-D	For pasture areas, remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment. Do not graze or feed treated wheat prior to maturity.
Rimfire	Forage = 30 days PHI, Grain and Straw = 71 days PHI.
Ro-Neet	No restrictions on the label.
Roundup Original MAX	Do not feed or graze treated grain sorghum fodder or ensile treated grain sorghum vegetation when using wiper applications. Do not feed or graze treated grain sorghum forage or fodder following applications with a hooded sprayer. Do not graze or harvest treated soybeans within 25 days of the last preharvest application (14 days if the application rate was 22 oz per acre or lower). Do not feed or graze sunflower forage following application.

Trade Name **Forage, Feed, and Grazing Restrictions**

Roundup PowerMAX	Do not feed or graze treated grain sorghum fodder or ensile treated grain sorghum vegetation when using wiper applications. Do not feed or graze treated grain sorghum forage or fodder following applications with a hooded sprayer. Do not graze or harvest treated soybeans within 25 days of the last preharvest application (14 days if the application rate was 22 oz per acre or lower). Do not feed or graze sunflower forage following application.
Roundup WeatherMAX	Do not feed or graze treated grain sorghum fodder or ensile treated grain sorghum vegetation when using wiper applications. Do not feed or graze treated grain sorghum forage or fodder following applications with a hooded sprayer. Do not graze or harvest treated soybeans within 25 days of the last preharvest application (14 days if the application rate was 22 oz per acre or lower). Do not feed or graze sunflower forage following application.
RT 3	Do not feed or graze treated grain sorghum fodder or ensile treated grain sorghum vegetation when using wiper applications. Do not feed or graze treated grain sorghum forage or fodder following applications with a hooded sprayer. Do not graze or harvest treated soybeans within 25 days of the last preharvest application (14 days if the application rate was 22 oz per acre or lower). Do not feed or graze sunflower forage following application.
Saber	Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied.
Salvo	Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied. Do not forage or feed corn fodder for 7 days following application.
Savage	Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied. Do not forage or feed corn fodder for 7 days following application.
Section	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
Select Max	Do not graze treated soybean fields or feed treated forage or hay to livestock. Do not apply within 15 days of grazing, feeding, or harvesting alfalfa for forage or hay.
Sencor	Treated soybean vines may be fed to livestock 40 days after application. Field corn treated with Sencor may be grazed or harvested for silage or grain 60 days after treatment. Do not graze or harvest alfalfa within 28 days after application.
Sequence	Do not graze or feed forage from treated corn for 30 days after application. Do not cut treated pod crops for hay within 120 days following application. Do not feed treated soybean (conventional) forage or hay for 30 days following application. Do not graze or feed treated forage or hay from Roundup Ready soybeans to livestock following a postemergence over the top application.
Shotgun	Do not graze or feed forage from treated areas for 60 days following application.
Silverado	Do not apply Silverado within 30 days of harvesting wheat forage, 50 days for hay, and 55 days for grain or straw.
Sinbar	No restrictions on the label.
sodium chlorate, others	
Sonalan	Do not graze or forage crop grown in treated soil or cut for hay or silage.
Sonic	Do not feed treated soybean forage or soybean hay to livestock.
Spartan	Do not feed treated soybean forage or soybean hay to livestock.
Spike	No restrictions on the label.
Starane	Do not allow livestock to graze treated areas or harvest treated forage within 47 days of application in corn, 40 days in grain sorghum, or 7 days in oats, barley or wheat. Do not apply less than 90 days before harvest of grain or stover in corn, 70 days in grain sorghum, or 40 days in oats, barley or wheat.
Starane + Saber	Do not allow livestock to graze treated areas or harvest treated forage within 14 days in oats, barley or wheat. Do not apply closer than 14 days before cutting of hay or harvesting of straw.
Starane + Salvo	Do not allow livestock to graze treated areas or harvest treated forage within 14 days in oats, barley or wheat. Do not apply closer than 14 days before cutting of hay or harvesting of straw.
Starane + Sword	Do not allow livestock to graze treated areas or harvest treated forage within 7 days in oats, barley or wheat. Do not apply closer than 40 days before cutting of hay or harvesting of straw.
Starane NXT	Do not allow livestock to graze treated areas within 45 days of application. Do not harvest hay or straw within 45 days of application.

Trade Name **Forage, Feed, and Grazing Restrictions**

Starane NXTcp	Do not allow livestock to graze treated areas within 45 days of application. Do not harvest hay or straw within 45 days of application.
Starane Ultra	Do not allow livestock to graze treated areas or harvest treated forage within 47 days of application in corn, 40 days in grain sorghum, or 7 days in oats, barley or wheat. Do not apply less than 90 days before harvest of grain or stover in corn, 70 days in grain sorghum, or 40 days in oats, barley or wheat.
Status	Do not harvest corn forage within 32 days of application.
Steadfast	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of application.
Steadfast ATZ	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of application.
Stealth	Livestock can graze or be fed forage from treated field corn or sorghum after 21 days following application. Do not feed lupine hay and forage or graze livestock in treated lupine fields. Do not feed or graze legume cover crops established following application. Livestock can graze or be fed forage from treated soybean fields. Do not feed forage or graze livestock in treated sunflower fields.
Sterling	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest.
Sterling Blue	CORN: May be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity. SORGHUM: Do not graze or feed treated sorghum forage or silage prior to mature grain stage. PASTURES and SMALL GRAINS/SORGHUM (if used for pasture or hay): Animals cannot be removed from treated area for slaughter prior to 30 days after last application; There is no waiting period between treatment and grazing for non-lactating animals; For lactating dairy animals, grazing and hay harvest restrictions are rate dependent. Up to 0.5 lb ai/A, allow 7 days before grazing and 37 days before hay harvest. Up to 1.0 lb ai/A, allow 21 days before grazing and 51 days before hay harvest. Up to 2.0 lb ai/A, allow 40 days before grazing and 70 days before hay harvest. SOYBEANS: Do not feed soybean fodder or hay following a preharvest application.
Sterling Plus	Corn can be harvested or grazed for feed once the crop has reached the ensilage (milk stage) or later in maturity. Sorghum may be harvested or grazed for feed once the crop has reached the mature grain stage.
Stinger	Do not permit lactating dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 1 week from treatment. Do not harvest hay from treated grain fields. Do not allow livestock to graze treated corn or harvest treated corn silage as feed within 40 days after last treatment. No restrictions on grazing or hay harvesting in rangeland or grass pasture situations. Do not feed or graze canola fields. Do not cut canola for forage. Do not graze sweet corn or popcorn fields for forage within 30 days of last treatment nor harvest for stover within 60 days of last treatment.
Storm	Do not use treated plants for feed or forage.
Stout	Do not graze or feed forage, hay or straw from treated areas to livestock within 30 days of Stout application.
SureStart	No restrictions on the label.
Surpass	No restrictions on the label.
Sword	Do not forage or graze dairy and meat animals on treated areas within seven days of slaughter.
Synchrony XP	Do not graze treated fields or harvest for forage or hay.
Targa	Do not graze livestock in treated areas. In addition, do not feed forage, hay, or straw from treated areas to livestock. Do not feed sugarbeet tops within 60 days of last application.
Topnotch	No restrictions on the label.
Tordon 22K	When applying more than 0.5 lb ai/A to grass pastures, do not cut grass for feed within two weeks after treatment. Meat animals grazing for up to two weeks after treatment should be removed from treated areas three days prior to slaughter. Do not graze lactating dairy animals on treated areas within two weeks after treatment. Do not graze or feed small grain forage from treated areas for 2 weeks after treatment. Do not harvest hay from treated grain fields.
Touchdown	Do not graze or feed corn forage or fodder following hooded sprayer applications. Allow a minimum of 50 days between postemergence application and harvest of forage of glyphosate-tolerant corn. Allow at least 25 days before grazing or harvesting for livestock feed following harvest aid applications in conventional soybeans. Do not graze or harvest treated glyphosate-tolerant soybeans for forage or hay. Do not graze or feed sunflower forage.

Trade Name	Forage, Feed, and Grazing Restrictions
-------------------	---

Treflan	Do not cut or graze alfalfa within 21 days after application.
trifluralin, generic	Do not cut or graze alfalfa within 21 days after application.
Trust	Do not cut or graze alfalfa within 21 days after application.
Ultra Blazer	Do not use treated plants for feed or forage.
Unity WDG	Do not graze or feed forage or grain from treated field corn to livestock within 30 days of application. Do not graze or feed forage or hay from treated areas to livestock (harvested straw may be used for bedding feed).
UpBeet	No restrictions on grazing or feeding of crop residue to livestock.
Valor	Do not graze treated fields or feed treated forage or hay to livestock.
Velpar	Do not graze or feed forage or hay to livestock within 30 days following application.
Velpar AlfaMax	Do not graze or feed forage to livestock within 30 days following application.
Velpar AlfaMax Gold	Do not graze or feed forage to livestock within 30 days following application.
Weedmaster	Remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals. Do not graze lactating dairy animals within 7 days of treatment. Treated grasses may be harvested for dry hay but do not harvest within 37 days of treatment.
WideMatch	Do not allow livestock to graze treated areas or harvest treated small grain forage within 7 days of application. Do not apply closer than 14 days before cutting small grains for hay or 40 days before harvesting of straw. Do not allow livestock to graze treated areas or harvest treated corn forage within 47 days of application. Do not apply less than 90 days before harvest of corn stover. Do not allow livestock to graze treated areas or harvest treated sweet corn forage within 31 days of application. Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.
WideMatch M	Do not allow livestock to graze treated areas or harvest treated small grain forage within 7 days of application. Do not apply closer than 14 days before cutting small grains for hay or 40 days before harvesting of straw. Do not allow livestock to graze treated areas or harvest treated corn forage within 47 days of application. Do not apply less than 90 days before harvest of corn stover. Do not allow livestock to graze treated areas or harvest treated sweet corn forage within 31 days of application. Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.
Yukon	Corn can be grazed or harvested for feed after the crop reaches the ensilage (milk) stage at least 30 days after application.