Weed control with AE F130360 in corn at Lamberton, MN in 2002. Getting, Jodie K. and Bruce D. Potter. The objective of this study was to evaluate AE F130360 tank-mix combinations for annual grass and annual broadleaf weed control in corn. This study was conducted on a Normania loam soil containing 4.2% organic matter, pH 6.5 and soil test P and K levels of 60 and 316 lb/A, respectively. A randomized complete block design with four replications and a plot size of 10 by 30 ft was used. The site was planted to oats in 2001 and was fall chiseled. The area was fertilized with 180 lb/A of nitrogen as urea. On May 2, 2002, Mycogen '4150LL' glufosinate resistant field corn was planted in 30-inch rows at a seeding rate of 33,000 seeds/A. Tefluthrin (Force 3G) was applied at 4 oz/1000 row feet in a T-band for the control of northern corn rootworm larvae. All treatments were applied with a tractor-mounted sprayer delivering 20 gpa at a pressure of 40 psi. The sprayer was equipped with 8002 flat-fan nozzles spaced 15 inches apart on the boom. Application dates, environmental conditions, plant sizes and rainfall data are listed below:

Date Treatment	May 3 PRE	June 5 POST
Temperature (F)		
air	48	68
soil (4 inch)	42	76
Relative humidity (%)	30	49
Wind (mph)	SE 10	calm
Sky	clear	clear
Soil moisture	dry	moist
Corn	<i>y</i>	
leaf no.	-	3-collar
height (inch)	-	4
Yellow foxtail		
leaf no.	-	1 to 4
height (inch)	-	1 to 3
no./ft ²	-	40
Common lambsquarters		
leaf no.	-	3 to 5
height (inch)	-	1 to 2
no./ft ²	-	4
Redroot pigweed		
leaf no.	-	3 to 5
height (inch)	-	1 to 2
no./ft ²	-	6
Rainfall after application ((inch)	
1 week	1.25	0.57
2 week	0.36	0.24
3 week	0.00	1.18

None of the herbicide treatments caused visible crop injury. On June 3, prior to POST treatments, flufenacet applied PRE gave 71% yellow foxtail control, 51% common lambsquarters control, and 59% redroot pigweed control. Isoxaflutole + atrazine and dimethenamid-P applied PRE provided 89% or greater control of yellow foxtail, common lambsquarters, and redroot pigweed. In September, the PRE/POST treatments gave 90 to 94% yellow foxtail control and 96% or greater common lambsquarters and redroot pigweed control. AE F130360 applied POST provided 81% yellow foxtail control. AE F130360 tank-mixed with [dicamba & SAN 1269H], mesotrione, or flufenacet gave 80 to 83%, 76 to 80%, and 81 to 84% control, respectively. All herbicide treatments resulted in 91% or greater common lambsquarters control and 89% or greater redroot pigweed control. (Southwest Research and Outreach Center, University of Minnesota, Lamberton).

Table. Weed control with AE F130360 in corn at Lamberton, MN in 2002 (Getting and Potte	Table. Weed control with AE F130360 in corn at	Lamberton, MN in 2002	(Getting and Potter)
---	--	-----------------------	----------------------

			SETLU		CHEAL				AMARE				
Treatment ^a Rate	Rate	6/3	6/21	6/28	9/9	6/3	6/21	6/28	9/9	6/3	6/21	6/28	9/9
	(lb/A or %)	(% control)											
POST (2 to 4-inch weeds)													
AE F130360+MSO+28%N	0.03+0.94%+2.5%	0	91	90	81	0	96	94	93	0	97	91	93
AE F130360+[Dica&SAN 1269H]	0.03+[0.064&0.0.026]	0	88	86	83	0	95	94	93	0	97	91	90
+MSO+28%N	+0.94%+2.5%												
AE F130360+[Dica&SAN 1269H]	0.03+[0.128&0.0.051]	0	87	85	80	0	97	94	93	0	98	93	91
+MSO+28%N	+0.94%+2.5%												
AE F130360+mesotrione+MSO+28%N	0.03+0.047+0.94%+2.5%	0	86	84	78	0	97	95	93	0	97	89	89
AE F130360+mesotrione+MSO+28%N	0.03+0.063+0.94%+2.5%	0	89	88	80	0	97	97	95	0	97	90	93
AE F130360+mesotrione+MSO+28%N	0.03+0.094+0.94%+2.5%	0	86	86	76	0	97	95	93	0	97	92	92
AE F130360+[Prim&Dica]	0.03+[0.019&0.1]	0	90	89	85	0	96	96	95	0	96	93	94
+MSO+28%N	+0.94%+2.5%												
AE F130360+flufenacet+MSO+28%N	0.03+0.23+0.94%+2.5%	0	91	89	84	0	92	93	91	0	97	95	92
AE F130360+flufenacet+MSO+28%N	0.03+0.3+0.94%+2.5%	0	88	88	81	0	98	97	96	0	98	98	97
[Nico&Rims]+mesotrione+Atra	[0.023&0.012]+0.063+0.25	0	91	90	84	0	97	97	95	0	97	96	95
+COC+AMS	+1%+2.0												
AE F130360+mesotrione+Atra	0.03+0.063+0.25	0	82	84	76	0	98	97	95	0	98	95	95
+COC+AMS	+1%+2.0												
Preemergence/POST (2 to 4-inch weeds)	1												
Isoxaflutole+Atra/	0.047+0.75/	89	94	95	93	93	98	98	97	93	98	98	97
AE F130360+MSO+28%N	0.03+0.94%+2.5%												
Dimt-P/[Dica&SAN 1269H]	0.94/[0.128&0.051]	93	92	94	94	89	98	98	96	94	97	98	96
+NIS+AMS	+0.25%+1.0												
Flufenacet/Gluf+Atra+AMS	0.375/0.31+0.5+3.0	71	94	98	90	51	98	98	96	59	98	96	96
<u>Checks</u>													
Weedy check		0	0	0	0	0	0	0	0	0	0	0	0
Weed-free		100	100	100	100	100	100	100	100	100	100	100	100
	LSD (0.10)	2.7	3.6	3.9	7.2	6.5	1.9	2.7	3.2	11.7	1.4	3.5	4.3

LSD (0.10) 2.7 3.6 3.9 7.2 6.5 1.9 2.7 3.2 11.7 1.4 3.5 4.3 ^a AE F130360 = Option 35 DF; Atra or atrazine = Aatrex 4L; [Dica&SAN 1269H] or [dicamba & SAN 1269H] = Distinct 70WG; Dimt-P or dimethenamid-P = Outlook 6L; flufenacet = Define 60DF; Gluf or glufosinate = Liberty 1.67L; isoxaflutole = Balance Pro 4L; mesotrione = Callisto 4L; [Nico&Rims] or [nicosulfuron & rimsulfuron] = Steadfast 75DF; [Prim&Dica] or [primsulfuron & dicamba] = Northstar 47.4WG; COC = crop oil concentrate; MSO = methylated seed oil; NIS = nonionic surfactant; 28%N = an aqueous solution of urea and ammonium nitrate; AMS = spray grade ammonium sulfate.